
Piton de la Fournaise Flank Displacement following the March 2007 eruption

Cayol Valerie^{*1}, Marine Tridon¹, Froger Jean-Luc¹, Keith Richards-Dinger², James Dieterich², and Aurélien Augier¹

¹Laboratoire Magmas et Volcans, Université Clermont-Auvergne – CNRS : UMR6524, Institut de recherche pour le développement [IRD] : UR163 – France

²Department of Earth Sciences, University of California, Riverside – United States

Abstract

The April 2007 eruption of Piton de la Fournaise was the biggest eruptive crisis of the 20th and 21st centuries for the volcano. InSAR captured a large (1.4 m) co-eruptive seaward slip of the volcano's eastern flank, which continued for more than a year at a decreasing rate after the end of the eruption. Co-eruptive uplift and post-eruptive subsidence were also observed. While it is generally agreed that volcano flank displacement might be induced by fault slip, we investigate whether this flank displacement might have been induced by a sheared sill, as suggested by observations of sheared sills at Piton des Neiges. To test this hypothesis, we invert a quadrangular curved source submitted to coeval pressure and shear stress changes. Post-eruptive displacement is well explained by slip and closure of a large fracture sub parallel to the topography (5 km by 8 km). The amount of closure is too large and the closure time too short to be explained by a thermally compacting sill, allowing to rule out the sill hypothesis. Co-eruptive displacement can be explained by a smaller (2 km by 2 km) fracture at the same location, submitted to a zero overpressure and a shear stress drop, which confirms that the determined structure is not a sill. We conclude that the fracture is a detachment fold, shallow enough to induce the observed coeval uplift. Observations are consistent with a rate weakening behavior for the patch which ruptured during the April 2007 eruption, while the creeping part of the fault is consistent with a rate strengthening behavior.

^{*}Speaker