

---

# Aseismic slip preceding the 2017 earthquakes sequence in the Mexican subduction zone.

Mathilde Radiguet<sup>\*1</sup>, Jorge Jara<sup>2</sup>, Ekaterina Kazachkina<sup>3</sup>, Louise Maubant<sup>1</sup>, Nathalie Cotte<sup>1</sup>, and Vladimir Kostoglodov<sup>3</sup>

<sup>1</sup>ISTerre, Institut des Sciences de la Terre – Université Grenoble Alpes – BP 53 - 38041 Grenoble cedex 9, France

<sup>2</sup>Laboratoire de Géologie – Ecole Normale Supérieure de Paris - ENS Paris – France

<sup>3</sup>Instituto de Geofísica – UNAM – Mexico DF, Mexico

## Abstract

The Mexican subduction zone is characterized by the occurrence of recurrent large slow slip events (SSEs), both in the Guerrero region, where the SSEs are among the largest observed worldwide, and in the Oaxaca region, where smaller, more frequent SSEs occur. In the past, complex interactions between aseismic processes and slow slip events have been identified in that region, in particular the triggering of two thrust earthquakes by ongoing SSEs [Graham et al. 2014, Radiguet et al. 2016] and the activation of SSEs and associated tremor signals by a distant earthquake [Zigone et al. 2012].

In this study, we investigate the aseismic-seismic interactions preceding the 2017 Mw 8.2 Tehuantepec and Mw 7.1 Puebla large intraplate earthquakes. The intraplate mechanisms of these earthquakes differ from previously observed aseismic-seismic interactions in the region, which consisted in the triggering of nearby thrust earthquakes. However, large scale aseismic slip (decoupling) have been proposed in Chile to explain interactions between deep and shallow seismicity [Jara et al. 2017].

We look for short-term and long-term variations in seismicity and surface displacements, by analyzing the SSN (Servicio Seismológico Nacional) seismicity catalog and the GPS time series (IG-UNAM and TLALOCNet).

At the time-scale of a few years preceding the 2017 earthquakes, we investigate the variations in the raw and background seismicity rate by declustering the seismicity catalog using an epidemic-type aftershock sequence (ETAS). Our results reveal an increase in the background seismicity rate in the Chiapas region, which initiates in early 2015, both in at shallow (< 30km) and intermediate (> 55km) depths. We complement this observation by an analysis of the surface deformations prior to the 2017 earthquakes. Several transient slip episodes (SSEs) are detected in the GPS time series, which locally modify the interseismic coupling. No significant long-term trend in the GPS time series, that could be associated with the observed variation in the seismicity rate, has been detected. However, at a shorter time scale, two slow slip events that initiate in May 2017, 4 months before the earthquakes, are identified along the Mexican subduction, both in Guerrero and Oaxaca.

References:

---

<sup>\*</sup>Speaker

Graham, S. E. et al. GPS constraints on the Mw7.5 Ometepec earthquake sequence, southern Mexico: coseismic and post-seismic deformation. *Geophys. J. Int.* 199, 200-218 (2014).

Jara, J., Socquet, A., Marsan, D., & Bouchon, M. (2017). Long-term interactions between intermediate depth and shallow seismicity in North Chile subduction zone. *Geophysical Research Letters*, 44.

D. Zigone, D. Rivet, **M. Radiguet**, M. Campillo, C. Voisin, N. Cotte, A. Walpersdorf, N. M. Shapiro, G. Cougoulat, P. Roux, V. Kostoglodov, A. Husker, and J. S. Payero (2012). Triggering of tremors and slow slip event in Guerrero, Mexico, by the 2010 Mw 8.8 Maule, Chile, earthquake. *J. Geophys. Res.*, 117(B9):B09304.

M. Radiguet, H. Perfettini, N. Cotte, A. Gualandi, B. Valette, V. Kostoglodov, T. Lhomme (2016). Triggering of the 2014Mw7.3 Papanaoa earthquake by a slow slip event in Guerrero, Mexico. *Nature Geoscience*, DOI : 10.1038/NGEO2817.